

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
28 December 2000 (28.12.2000)

PCT

(10) International Publication Number  
**WO 00/78306 A1**

(51) International Patent Classification<sup>7</sup>: **A61K 31/28**

J. [US/US]; 3220 S. Old Ridge Road, Columbia, MO 65203 (US).

(21) International Application Number: PCT/US00/17341

(22) International Filing Date: 23 June 2000 (23.06.2000)

(74) Agents: **KOHN, Kenneth, I.** et al.; Kohn & Associates, Suite 410, 30500 Northwestern Highway, Farmington Hills, MI 48334 (US).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/140,576 23 June 1999 (23.06.1999) US  
60/156,151 27 September 1999 (27.09.1999) US

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(71) Applicant (*for all designated States except US*): **THE CURATORS OF THE UNIVERSITY OF MISSOURI** [US/US]; 615 Locust Street Building, 3rd floor, Columbia, MO 65211-1400 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventor: **KAMEI, Hideo** (deceased).

Published:

— With international search report.

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **KATTI, Kattesh, V.** [IN/IN]; 3901 Woods Edge Road, Columbia, MO 65203 (US). **VOLKERT, Wynn, A.** [US/US]; 2808 Butternut Ct., Columbia, MO 65203 (US). **HOFFMAN, Timothy,**

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: GOLD-CONTAINING CHEMOTHERAPEUTIC AGENTS

(57) Abstract: There is provided a complex for use as a therapeutic pharmaceutical, the complex has a ligand containing at least one hydroxyalkyl phosphine donor group bound to a gold atom to form a stable gold-ligand complex. Also provided is a method of treating cancer by administering an effective amount of a complex having a ligand of at least one hydroxyalkyl phosphine group bound to a gold atom to form a stable gold-ligand complex. Also provided is a method of preventing the metastasis of cancer and arresting cell growth by administering an effective amount of a complex having a ligand of at least one hydroxyalkyl phosphine group bound to a gold atom to form a stable gold-ligand complex.



**WO 00/78306 A1**

## GOLD-CONTAINING CHEMOTHERAPEUTIC AGENTS

### Technical Field

5

The present invention relates to pharmaceuticals for use as therapeutic agents. More specifically, the present invention relates to pharmaceuticals containing therein gold for use as therapeutic agents.

10

### Background of the Invention

Gold, in elemental form, has been employed for centuries as an antipruritic to relieve the itching palm. In more modern times, the observation by Robert Koch in 1890 that gold inhibited *Mycobacterium tuberculosis in vitro* led to trials in arthritis and lupus erythematosus, thought by some to be tuberculous manifestations. Later observations of success in treating chronic arthritis stimulated interest in gold therapy (chrysotherapy). At present, gold is employed in the treatment of rheumatoid arthritis; usually it is reserved for patients with progressive disease who do not obtain satisfactory relief from therapy with NSAIDs. However, gold compounds are among the agents that are used in an attempt to arrest the progress of the disease and to induce remissions; these are sometimes called disease-modifying drugs, although this is probably a misnomer (Edmonds et al., 1993). Since degenerative lesions do not regress once formed, there is an increasing tendency to attempt to induce remission early in the course of the disease. Such therapy is often initiated with gold, which although potentially beneficial, causes a high incidence of toxicity (Felson et al., 1992; Cash and Klippel, 1994).

Gold compounds can suppress or prevent, but not cure, experimental arthritis and synovitis due to a number of infectious and chemical agents.

Gold compounds have minimal antiinflammatory effects in other circumstances and cause only a gradual reduction of the signs and symptoms of inflammation associated with rheumatoid arthritis. Although many effects of these drugs have been observed, which, if any, are related to the therapeutic effects of gold in rheumatoid arthritis is unknown. Perhaps the best hypotheses relate to the capacity of gold compounds to inhibit the maturation and function of mononuclear phagocytes and of T cells, thereby suppressing immune responsiveness. Decreased concentrations of rheumatoid factor and immunoglobulins often are observed in patients who are treated with gold.

In experimental animals, gold is sequestered in organs that are rich in mononuclear phagocytes, and it selectively accumulates in the lysosomes of type A synovial cells and other macrophages within the inflamed synovium of patients who are treated with gold compounds. Moreover, the administration of gold thiomalate to animals depresses the migration and phagocytic activity of macrophages in inflammatory exudates, and chrysotherapy reduces the augmented phagocytic capacity of blood monocytes from patients with rheumatoid arthritis. Other mechanisms of action of gold compounds have been suggested, but none is generally accepted. These include inhibition of prostaglandin synthesis, interference with complement activation, cross-linking of collagen, and inhibition of the activity of lysosomal and other enzymes, including protein kinase C, in T cells.

Considerable research has been focused on the development of water-soluble gold-containing compounds because of their potential in medical applications [1]. The first application of gold-containing compounds came from their use in rheumatoid arthritis. The compounds used in the treatment of rheumatoid arthritis were aurothiomalate (Myocrisin) and aurothioglucose (Solgano) as depicted in Figure 1. In 1985, another gold containing

compound, auranofin, [(2,3,4,6-tetra-O-aceetyl-1-thio- $\beta$ -D-glucopyranosato-S)-(triethylphosphine) gold(I)] was shown to be effective for the treatment of rheumatoid arthritis [2,3]. Several studies have demonstrated that this agent is superior to the traditional chrysotherapeutic drugs. Auranofin and related gold(I) compounds have been found to be active against interperitoneal P388 leukemia and are also cytotoxic to specific tumor cells [4,5]. Mirabelli et al. screened the  $\mu$ -[bis(diphenylphosphino)ethane] digold complex [dppe(AuCl)<sub>2</sub>] (Figure 2) for antitumor activities [6]. Such digold complexes rearranged to give tetrahedral complexes of the type [Au(dppe)<sub>2</sub>].

10

The tetrahedral arrangement of ligands around gold (as in Au(dppe)<sub>2</sub>; Figure 2) allowed better stabilization of the metal center through chelate effects. Such tetrahedrally-bound phosphine ligands around the gold center are more inert to substitution by potential thiolate ligands that could be encountered in a biological environment.

15

It was suggested that the mechanism of action for [Au(dppe)<sub>2</sub>]Cl was the formation of DNA protein cross-links [7,8]. The lack of affinity of gold(I) for oxygen and nitrogen containing ligands resulted in poor reactivity with the bases of DNA.

20

The gold compound [Au(dppe)<sub>2</sub>]Cl demonstrated marked activity against peritoneal cancer cells. However, this compound was found to be only slightly active against solid tumor models. This compound could not proceed to clinical trials due to problems with cardiotoxicity as revealed in preclinical toxicology studies [9].

25

Failure to identify an effective gold-containing antitumor agent stems from the difficulties associated with the development of gold compounds with optimum hydrophilicity/lipophilicity, toxicity and activity toward specific tumors.

30

Therefore, an improved understanding of the molecular and biochemical mechanism of gold compounds can provide the impetus for new advances in the antitumor applications of gold compounds. It would also be useful to develop an antitumor gold compound which is not toxic to the patient.

5

It would therefore be useful to develop pharmaceuticals to develop stable non-radioactive gold complexes for use as chemotherapeutic agents.

### **Summary of the Invention**

10

According to the present invention, there is provided a complex for use as a therapeutic pharmaceutical, the complex including a ligand containing at least one hydroxyalkyl phosphine donor group bound to a gold atom to form a stable gold-ligand complex. Also provided is a method of treating cancer by administering an effective amount of a complex having a ligand of at least one hydroxyalkyl phosphine group bound to a gold atom to form a stable gold-ligand complex. Also provided is a method of preventing the metastasis of cancer and arresting cell growth by administering an effective amount of a complex having a ligand of at least one hydroxyalkyl phosphine group bound to a gold atom to form a stable gold-ligand complex.

15  
20

### **Brief Description of the Drawings**

Other advantages of the present invention will be readily appreciated as the same becomes better understood by reference to the following detailed description when considered in connection with the accompanying drawings wherein:

25

Figure 1 illustrates the prior art gold-containing compound;

Figure 2 illustrates further prior art gold-containing complexes;

Figure 3 illustrates representative hydroxymethyl phosphine (HMP  
5 ligands);

Figure 4 illustrates the synthesis of trihydroxymethyl phosphine gold complexes;

10 Figure 5 illustrates variations of alkyl chain length of hydroxymethyl phosphines; and

Figure 6 illustrates alternative approaches of chelating HMP groups to produce tetrahedral gold compounds.

15

#### **Detailed Description of the Preferred Embodiment**


Generally, the present invention provides a complex for use as a therapeutic pharmaceutical. The complex includes a ligand including at least  
20 one hydroxyalkyl phosphine donor group bound to a non-radioactive gold atom to form a gold ligand complex that is stable. That is, the invention provides a ligand system containing at least one hydroxyalkyl phosphine donor group for use in forming complexes with non-radioactive gold metals wherein the complexes have high *in vitro* and/or *in vivo* stability.

25

The phosphorous ligands were chosen since the phosphorous atom provides a plethora of electron density that promotes formation of highly stable ligand metal bonds. This can occur even with non-radioactive gold metal in its higher oxidation states.

The hydroxyalkyl phosphine ligand is complexed with a non-radioactive gold metal, generally a gold (I) compound. These complexes contain a ratio of ligand-to-gold that is greater than or equal to 1:1 which makes the resulting  
 5 chelates small and well-defined.

A complex according to the present invention can include a complex of the formula




10

wherein M is non-radioactive gold metal in a reduced oxidation state, n is 1-6, and A is an alkyl group. In a preferred embodiment of the present invention, A is  $-CH_2-$ . Additionally, A can include  $-C_2H_4-$  and iso- or normal- $C_3H_6-$ .

15

The non-radioactive gold-ligand complexes can include other donor atoms or groups on the same ligand as the donor hydroxyalkyl phosphine group. These other donor groups can include N, S, O, or P atoms for coordinating the non-radioactive gold atom. In addition, the donor groups can further include amines, amides, thiols, carboxyls, and hydroxyls for  
 20 coordinating the non-radioactive gold atom.

In another preferred embodiment of the present invention, the complexes can include a bidentate ligand of the formula




25

wherein A is  $-CH_2-$ ,  $-C_2H_4-$ , or iso- or normal-  $C_3H_6$ , and X is  $-(CH_2)_n$  - where  $n=1-4$ ,  $-CH_2CHR-$ ,  $-CH_2CHRCH_2-$ ,  $-CHRCH_2CH_2-$ , R'-aromatic where R' is H, an alkyl group of  $C_1-C_4$ , an aromatic group,  $-OH$ ,  $-SH$ ,  $-NH_2$ ,  $-COOH$ , activated esters, N-hydroxysuccinimides, benzyl isothiocyanate, alkyl halides, or cyclohexyldiimide. Specific examples of bidentate ligands used to complex  
 30 with gold can include 1,2-bis (bis(hydroxymethyl)phosphino)benzene (HMPB,


1) and 1,2-bis(bis(hydroxymethyl)phosphino)ethane (HMPE, 2) as set forth below. The formation of non-radioactive gold complexes according to the present invention with the ligands HMPB and HMPE are shown.

5 In further preferred embodiments, complexes according to the present invention are contemplated which include multidentate ligands of the formula



wherein A is  $-\text{CH}_2-$ ,  $-\text{C}_2\text{H}_4-$ , or iso- or normal- $\text{C}_3\text{H}_6-$ , and X is  $-(\text{CH}_2)_n-$  where  $n=1-4$ ,  $-\text{CH}_2\text{CHR}-$ ,  $-\text{CH}_2\text{CHRCH}_2-$ ,  $-\text{CHRCH}_2\text{CH}_2-$ , or  $\text{R}'$ -aromatic where  $\text{R}'$  is H, an alkyl group of  $\text{C}_1-\text{C}_4$ , an aromatic group,  $-\text{OH}$ ,  $-\text{SH}$ ,  $-\text{NH}_2$ ,  $-\text{COOH}$ , activated esters, N-hydroxysuccinimides, benzyl isothiocyanate, alkyl halides, or cyclohexyldiimide, and Y is  $\text{CH}_2-$ ,  $-\text{C}_2\text{H}_4-$  or  $-\text{C}_3\text{H}_6-$ . Along the lines of this embodiment, further embodiments can exist wherein all of the donor atoms can be phosphorous atoms. Additionally, embodiments are contemplated  
15 wherein at least one donor group is a hydroxyalkyl phosphine group.


Furthermore, complexes according to the present invention are contemplated wherein two donor atoms are hydroxyalkyl phosphine phosphorous-atoms and two donor atoms are atoms other than phosphorous-atoms. These complexes have the general formula  
20



wherein A is  $-\text{CH}_2-$ ,  $-(\text{CH}_2)_2-$ , or iso- or normal- $\text{C}_3\text{H}_6-$ , K includes donor atoms or groups selected from the group consisting of  $-\text{N}-$ ,  $-\text{N}(\text{R})^+-$ ,  $-\text{N}(\text{H})-$ , Ag, and  $-\text{S}-$ , Y is  $-\text{CH}_2-$ ,  $-(\text{CH}_2)_2-$ , or iso- or normal- $\text{C}_3\text{H}_6-$ , X is  $-\text{CH}_2-$ ,  $-(\text{CH}_2)_2-$ ,  $-(\text{CH}_2)_3-$ ,  $-(\text{CH}_2)_4-$ ,  $-\text{CH}_2\text{CHR}-$ ,  $-\text{CH}_2\text{CHRCH}_2-$ ,  $-\text{CHRCH}_2\text{CH}_2-$ , or  $\text{R}'$ -aromatic wherein  $\text{R}'$  and R can be the same or different and are selected from H,  $-\text{OH}$ ,  $-\text{SH}$ ,  $-\text{NH}_2$ ,  $-\text{COOH}$ , activated esters, N-hydroxysuccinimides, benzyl isothiocyanate, alkyl halides, and cyclohexyldiimide.  
25



A variant of this embodiment can include a complex wherein two donor atoms are hydroxyalkyl phosphine phosphorous-atoms and two donor atoms are nitrogen-atoms (P2N2). These complexes can have the general formula




- 5 wherein X is  $-CH_2-$ ,  $-(CH_2)_2-$ ,  $-C_3H_6-$ , A is  $-CH_2-$ ,  $-(CH_2)_2-$ ,  $-C_3H_6-$ , E is O or S. R can be the same or different and is selected from H,  $-OH$ ,  $-SH$ ,  $-NH_2$ ,  $-COOH$ , activated esters, N-hydroxysuccinimides, benzyl isothiocyanate, alkyl halides, or cyclohexyl diimide.

- 10 N is nitrogen, and Y is  $-CH_2-$ ,  $-(CH_2)_2-$ , or iso- or normal- $C_3H_6-$ .

An additional embodiment is contemplated wherein two donor atoms are hydroxyalkyl phosphine phosphorous-atoms and two donor atoms are sulfur-atoms (P2S2).

15

Complexes contemplated under this embodiment have the general formula



- 20 wherein X is  $-CH_2-$ ,  $-(CH_2)_2-$ ,  $-C_3H_6-$ , A is  $-CH_2-$ ,  $-(CH_2)_2-$ ,  $-C_3H_6-$ , E is O or S. R can be the same or different and is selected from H,  $-OH$ ,  $-SH$ ,  $-NH_2$ ,  $-COOH$ , activated esters, N-hydroxysuccinimides, benzyl isothiocyanate, alkyl halides, or cyclohexyldiimide, S is sulfur, and Y is  $-CH_2-$ ,  $-(CH_2)_2-$ , or iso- or normal- $C_3H_6-$ .

- 25 A "therapeutically effective amount" is an amount of a complex of the present invention that, when administered to a patient, ameliorates a symptom of the specific disease or condition being treated. A therapeutically effective amount of a complex of the present invention can easily be determined by one skilled in the art by administering a quantity of a complex
- 30 to a patient and observing the result. In addition, those skilled in the art are

familiar with identifying patients having the particular disease or condition and are readily able to identify patients who suffer from these diseases or conditions.

- 5           The complexes of the present invention can be administered to a patient alone or as part of a composition that contains other components such as excipients, diluents, and carriers, all of which are well-known in the art. The compositions can be administered to humans and animals either orally, rectally, parenterally (intravenous, by intramuscularly or subcutaneously),  
10   intracisternally, intravaginally, intraperitoneally, intravesically, locally (powders, ointments or drops), or as a buccal or nasal spray.

- Compositions suitable for parenteral injection can comprise physiologically acceptable sterile aqueous or nonaqueous solutions,  
15   dispersions, suspensions or emulsions, and sterile powders for reconstitution into sterile injectable solutions or dispersions. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (propyleneglycol, polyethyleneglycol, glycerol, and the like), suitable mixtures thereof, vegetable oils (such as olive oil) and injectable  
20   organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersions and by the use of surfactants.

- These compositions can also contain adjuvants such as preserving,  
25   wetting, emulsifying, and dispensing agents. Prevention of the action of microorganisms can be ensured by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, and the like. It may also be desirable to include isotonic agents, for example, sugars, sodium chloride, and the like. Prolonged absorption of the injectable

pharmaceutical form can be brought about by the use of agents delaying absorption, for example, aluminum monostearate and gelatin.

Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active complex is admixed with at least one customary inert excipient (or carrier) such as sodium citrate or dicalcium phosphate or (a) fillers or extenders, as for example, starches, lactose, sucrose, glucose, mannitol, and silicic acid; (b) binders, as for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose and acacia; (c) humectants, as for example, glycerol; (d) disintegrating agents, as for example, agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain complex silicates and sodium carbonate; (e) solution retarders, as for example paraffin; (f) absorption accelerators, as for example, quaternary ammonium complexes; (g) wetting agents, as for example, acetyl alcohol and glycerol monostearate; (h) adsorbents, as for example, kaolin and bentonite; and (i) lubricants, as for example, talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, or mixtures thereof. In the case of capsules, tablets, and pills, the dosage forms may also comprise buffering agents.

20

Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols, and the like.

Solid dosage forms such as tablets, capsules, pills, and granules can be prepared with coatings and shells, such as enteric coatings and others well-known in the art. They may contain pacifying agents and can also be of such composition that they release the active complex or complexes in a certain part of the intestinal tract in a delayed manner. Examples of embedding compositions which can be used are polymeric substances and

30

waxes. The active complexes can also be in micro-encapsulated form, if appropriate, with one or more of the above-mentioned excipients.

Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs. In addition to the active complexes, the liquid dosage forms can contain inert diluents commonly used in the art, such as water or other solvents, solubilizing agents and emulsifiers, as for example, ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1.3-butylene glycol, dimethylformamide, oils, in particular, cottonseed oil, groundnut oil, corn germ oil, olive oil, castor oil and sesame oil, glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan or mixtures of these substances, and the like.

Besides such inert diluents, the composition can also include adjuvants, such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

Suspensions, in addition to the active complexes, can contain suspending agents, as for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, or mixtures of these substances, and the like.

Compositions for rectal administrations are preferably suppositories which can be prepared by mixing the complexes of the present invention with suitable nonirritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax, which are solid at ordinary temperatures but liquid at body temperature and therefore, melt in the rectum or vaginal cavity and release the active component.

Dosage forms for topical administration of a complex of this invention include ointments, powders, sprays and inhalants. The active component is admixed under sterile conditions with a physiologically acceptable carrier and any preservative, buffers, or propellants as may be required. Ophthalmic formulations, eye ointments, powders, and solutions are also contemplated as being within the scope of this invention.

The complexes and/or compositions of the present invention can be administered to a patient at dosage levels in the range of about 0.1 to about 1,000 mg per day. For a normal human adult having a body weight of about 70 kilograms, a dosage in the range of about 0.01 to about 7000 mg per kilogram of body weight per day is sufficient. The specific dosage used, however, can vary. For example, the dosage can depend on a number of factors including the requirements of the patient, the severity of the condition being treated, and the pharmacological activity of the complex being used. The determination of optimum dosages for a particular patient is well-known to those skilled in the art.

In addition, the complexes of the present invention can exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of the present invention.

25

The complex of the present invention can be coadministered with an additional therapeutic agent. This therapeutic can include, but is not limited to, chemotherapeutic agents. Preferably, the complex of the present invention and the coadministered therapeutic agent work in conjunction with one another to create a more sustained effect. These two therapeutic agents

30

can be either administered in one pharmaceutically acceptable carrier or separately.

When the non-radioactive analogue of the gold complex of THP  
5 [(Au(THP)<sub>4</sub>)] (Figure 4) was tested for tumor cell suppression against specific human cancer cells lines, it came as a total surprise that this compound exhibited remarkable activity in suppressing tumor growth in *in vitro* conditions. The data for 50% and 70% tumor growth suppression of cells derived from human colon carcinoma are summarized in Tables 1 and 2.

10 The data in Table 1 demonstrates that 50% tumor growth suppression occurs at ~2μg of the gold compound. This invention is significant in that it demonstrates high efficacy of tumor growth suppression under extremely low doses of the new antitumor agent (Au(THP)<sub>4</sub>]Cl. The tumor growth  
15 suppression for cells derived from human gastric carcinoma have been tested. The data for 50% and 70% tumor growth suppression are summarized in Tables 2 and 4 respectively. This 50% tumor growth suppression of cells derived from human gastric carcinoma occurs at about 10mg of [Au(THP)<sub>4</sub>]Cl. This is the first demonstration that hydroxymethyl  
20 phosphine (HMP)-bound gold compounds display high efficacy in suppressing tumor growth of specific cells derived from human carcinoma.

Clinically, cisplatin is widely used as a chemotherapeutic agent in the treatment of human cancer. Use of cisplatin is generally associated with  
25 severe toxic side effects that include decrease in blood cell numbers, kidney dysfunction, etc. Therefore, cisplatin cannot be used in treating cancer patients for longer periods of time. Gold-containing compounds are less toxic and their non-toxic dose is generally higher than that of cisplatin. Therefore, it is practical to treat cancer patients using gold-containing chemotherapeutic  
30 agents for longer durations. In this context, the tumor growth suppression

data reported for  $[\text{Au}(\text{THP})_4]\text{Cl}$  demonstrates the potential of using this (and related) new generation of gold-containing compounds in treating cancer bearing patients.

5           The above discussion provides a factual basis for the use of gold-containing chemotherapeutic agents. The methods used with and the utility of the present invention can be shown by the following non-limiting examples and accompanying figures.

## 10       **EXAMPLES**

### **Example 1**

#### **Histogram Studies:**

15

Histogram studies were carried out using HCT-15 cells. In the enclosed figure that summarize histogram results the cells are numbered as follows: HCT-80 is the control; HCT-81 refers to a sample with 200  $\mu\text{g}/\text{ml}$  of the gold compound; HCT-82 refers to a sample with 40  $\mu\text{g}/\text{ml}$  of gold  
20       compound.

25

HCT-15 cells were incubated in 5%  $\text{CO}_2$  incubator for two days. Then the cells were harvested and analyzed using fluorescent techniques. As shown on the enclosed figures, the percent of G1 phase in the cell cycle is significantly less for the control as compared to the cell samples that contained 200 or 40  $\mu\text{g}$  of gold compound. This is an important finding because it clearly demonstrates that the gold compound elongates G1 phase and consequently slows cell growth. This finding is in sharp contrast to the cytotoxic side effects of cisplatin which is known to cause disassociation of  
30       many phases. Additionally, most agents which are cytotoxic, block S or M+

G2 phase after culturing for two days. The mode of cell suppression via G1 phase elongation, as observed for the gold compound, therefore functions differently.

5     ***In vivo* studies:**

          The data of *in vivo* studies are summarized in Table: In vivo below. As shown in this table, the gold compound causes remarkable survival of tumor bearing mice. Administration of gold compound in doses significantly larger  
10    than the amount shown in this table (20-25 µg/kg) did not result in deaths of tumor bearing mice. Cisplatin at such doses resulted in the death of animals.

**Side effects:**

15           Preliminary findings, to date, have indicated no toxic cardiovascular or nephrotoxic side effects (in mice models) for the {Au[THP]<sub>4</sub>C1} compound.



**TABLE: *In vivo***  
**Results of *in vivo* study of [Au(THP)<sub>4</sub>Cl]**

5                    **Male 3.5 weeks weighing 17 to 20 g. mice.**  
**Balb/C mice inoculated 30,000 of Meth/A cells i.p. on 6/9/99**

		mice administered with Au compound of 1 mg/kg s.c. for three times daily for three days only (3 mg/kg total) (n=17)	mice administered with Au compound of 0.2 mg/kg s.c. for three times daily for three days only (0.6 mg/kg total)(n=15)
10			
15	<b>Control mice (n=17) date</b>		
20	6/9 0/17	0/17	0/15
	6/25 1/16	0/17	0/15
	6/26 1/15	0/17	0/15
	6/28 1/14	0/17	0/14
	6/29 1/13	0/17	0/13
25	6/30 0/13	0/17	0/13
	7/1 0/13	0/17	0/13
	7/2 3/10	1/16	1/12
	7/3 2/8	0/16	1/11
	7/5 0/8	1/15	0/11
30	7/6 0/8	0/15	1/10
	7/8 0/8	0/15	0/10
	7/9 0/8	0/15	0/10
	7/10 0/8	0/15	0/10
	7/14 0/8	0/15	0/10
35	7/20 0/2	0/15	0/10

a/b: a=number of dead mice      b=number of living mice

40                    The data summarized in Tables 1-4 demonstrate the efficacy of  
[Au(THP)<sub>4</sub>]Cl in suppressing tumor growth of human colon cancer and human

gastrin cancer cells. This compound is effective in treating various other carcinoma such as prostate cancer, breast cancer, brain tumors and also pancreatic cancer. The pharmacokinetic and drug action of Au-containing chemotherapeutic agents can be readily altered by systematic modifications of HMP ligands. One way is to vary the alkyl chains of alkyl hydroxy groups as shown in Figure 5. The other approach would involve utility of chelating HMP compounds such as HMPB and HMPE to produce tetrahedral gold compounds as shown in Figure 6. Therefore, chemical modifications of ligand backbones allows systematic tuning of gold pharmacophore characteristics. This aspect is important in terms of the design and development of gold-containing chemotherapeutic agents with optimum hypophilic/lipophilic characteristics and charge on the gold center that may be needed in the treatment of specific human carcinoma.

Potential applications of  $[\text{Au}(\text{THP})_4]\text{Cl}$  and related analogues include (but are not limited to): (i) treatment of small cell lung cancer; (ii) treatment of prostate cancer; (iii) treatment of breast, colon, pancreatic cancers and other kinds of malignancies; and (iv) treatment of rheumatoid arthritis.

Throughout this application, various publications, are referenced by author and year. Full citations for the publications are listed below. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention pertains.

25

The invention has been described in an illustrative manner, and it is to be understood that the terminology which has been used is intended to be in the nature of words of description rather than of limitation.

Obviously, many modifications and variations of the present invention are possible in light of the above teachings. It is, therefore, to be understood that within the scope of the appended claims, the invention may be practiced otherwise than as specifically described.

## REFERENCES

1. Kamei, H.; Koide, T.; Kojima, T.; Hashimoto, Y.; Hasegawa, M.  
Cancer Biotherapy & Radiopharmaceuticas. 1998, 13, 403.
- 5 2. Fricker, S.P. Transition Met. Chem. 1996, 21, 377.
3. Berglof, F.E.; Berglof, K.; Walz, D.T. J. Rheumatol. 1978, 5, 68.
4. Simon, T.M.; Kunishima, D.H.; Vibert, G.J.; Lorber, A. Cancer Res.  
1981, 41, 94.
5. Mirabelli, C.K.; Johnson, R.K.; Sung, C.M.; Faucette, L.; Muirhead, K.;  
10 Crooke, S.T. Cancer Res. 1985, 45, 32.
6. Snyder, R.M.; Mirabelli, C.K.; Johnson, R.K.; et al. Cancer Res. 1986,  
46, 5064.
7. Berners, S.J.; Jarrell, P.S.; Sadler, P.J. Inorg. Chem. 1987, 26, 3074.
8. Sadler, P. Agnew, Chem. Int. Ed. 1999, 38, 1512.
- 15 9. Hoke, G.D.; Macia, R.A.; Meunier, P.J.; Bugelski, P.J.; Mirabelli, C.K.;  
Rush, G.F.; Matthews, W.D. Toxicol. Appl. Pharmacol. 1989, 100, 293.
10. Volkert, W.A.; Goeckeler, W.F.; Ehrhardt, G.J.; Ketrin, A.R. J. Nucl.  
Med. 1991, 32, 174.
11. Schubiger, P.A.; Alberto, R.; Smith, A. Bioconjugate Chem. 1996, 7,  
20 165.
12. Shi, J.C.; Huang, X.Y.; Wu, D.X.; Liu, Q.T.; Kang, B.S. Inorg. Chem.  
1996, 35, 2742.
13. Rush, G.F.; Alberts, D.W.; Meunier, D.; Leffler, K.; Smith, P.F.  
Toxicologist 1987, 7, 59.
- 25 14. Katti, K.V. Curr. Sci. 1996, 70, 219.
15. (a) Reddy, V.S.; Berning, D.E.; Katti, K.V.; Barnes, C.L.; Volkert, W.A.;  
Ketrin, A.R. Inorg. Chem. 1996, 35, 1753. (b) Reddy, V.S.; Katti, K.V.;  
Barnes, C.L. J. Chem. Soc. Dalton Trans. 1996, 1301. (c) Reddy, V.S.; Katti,  
K.V.; Barnes, C.L. Inorg. Chem. Acta. 1995, 240, 367.
- 30 16. Berning, D.E.; Katti, K.V.; Sing, P.R.; Kiggenbotham, C.; Reddy, V.S.;  
Volkert, W.A. Nucl. Med. Biol. 1996, 23, 617.

17. Smith, C.J.; Katti, K.V. et al. Nucl. Med. Biol. 1997, 24, 685.
18. Berning, D.E.; Katti, K.V. et al. Nucl. Med. Biol. 1998, 25, 577.

## CLAIMS

What is claimed is:

- 5           1.       A complex for use as a therapeutic pharmaceutical, said complex comprising:  
              a ligand comprising at least one hydroxyalkyl phosphine donor group bound to a non-radioactive gold atom to form a stable gold-ligand complex.
- 10           2.       The complex according to claim 1, wherein said gold atom is a gold(I) compound.
3.       The complex according to claim 1 wherein said gold-ligand complex further includes a pharmaceutically acceptable carrier.
- 15           4.       The complex according to claim 1, wherein said gold-ligand complex further includes therapeutic agent.
5.       A method of treating cancer comprising the steps of:  
20           administering an effective amount of a complex comprising a ligand comprising at least one hydroxyalkyl phosphine group bound to a non-radioactive gold atom to form a stable gold-ligand complex.
6.       A method of preventing the metastasis of cancer comprising the  
25           steps of:  
              administering an effective amount of a complex comprising a ligand comprising at least one hydroxyalkyl phosphine group bound to a non-radioactive gold atom to form a stable gold-ligand complex.

7. A method of arresting cell growth comprising the steps of:  
administering an effective amount of a complex comprising a ligand  
comprising at least one hydroxyalkyl phosphine group bound to a non-  
5 radioactive gold atom to form a stable gold-ligand complex.









Figure - 1




Figure - 2


## REPRESENTATIVE HYDROXYMETHYL PHOSPHINE(HMP) LIGANDS


Figure - 3

## SYNTHESIS OF TRIHYDROXYMETHYL PHOSPHINE GOLD COMPLEXES

Figure - 4

## VARIATIONS OF ALKYL CHAIN LENGTHS OF HYDROXYMETHYL PHOSPHINES

Figure - 5

**Figure - 6**

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US00/17341

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/28

US CL : 514/108, 495

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/108, 495

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5,843,993 A (KATTI et al.) 01 December 1998, see entire document	1-7

☐

Further documents are listed in the continuation of Box C.

☐

See patent family annex.

\*

Special categories of cited documents:

"A"

document defining the general state of the art which is not considered to be of particular relevance

"E"

earlier document published on or after the international filing date

"L"

document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O"

document referring to an oral disclosure, use, exhibition or other means

"P"

document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&"

document member of the same patent family

Date of the actual completion of the international search

07 SEPTEMBER 2000

Date of mailing of the international search report

03 OCT 2000

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

*Frederick Kraus*  
FREDERICK KRAUS

Telephone No. (703) 308-1235